

---

## Letter to the Editor

---

### NUCLEAR DNA CONTENT AND GENOME SIZE OF TROUT AND HUMAN

In their recent paper, Thomas et al. (1) discussed design, resolution, sensitivity, and reproducibility of the National Aeronautics and Space Administration/American Cancer Society flow cytometer. The results of their study demonstrated high stability and sensitivity of the instrument, which is suitable for detection of near-diploid tumor cells. Although the performance of the instrument is impressive, we believe that Thomas et al. made serious errors in calculating the DNA contents of trout and human.

To estimate the DNA content of human cells in picograms of DNA, the authors used trout red blood cell nuclei as the internal reference standard and assumed 2.37 pg of DNA for a trout nucleus. With this value, the DNA contents of human female and male nuclei were estimated to be equal to 3.77 and 3.70 pg of DNA, respectively. The new estimates for trout and human differ drastically from previous ones, which range from 4.9 to 6.3 pg for various species of trout (2-5) and from 6.0 to 7.0 pg for human (5-9). Because trout and human nuclei are often used as internal reference standards to determine genome size in animals and plants (5,10,11), the results published by Thomas et al. (1) need correction to avoid serious mistakes.

Careful reading of the paper showed that the DNA content of trout (the authors did not specify the species) was derived after determining the ratio of mean DNA content of human male to trout to be 1.565. The DNA amount of the human nucleus, 3.70 pg, was calculated with the assumption that there are  $6.162 \times 10^9$  nucleotides for the human male nucleus and that a mean nucleotide molecular weight of 360 g/mol. We believe these data are not correct.

The ratio of human to trout DNA content determined by Thomas et al. (1) differs significantly from that of other reports. For instance, Vindeløv et al. (12) found that rainbow trout has 80% of human DNA content.

The most recent estimate of the size of the human diploid male genome is  $6.294 \times 10^9$  nucleotide pairs (13).

Mean nucleotide molecular weight is not 360 (Table 1).

The amount of DNA in a human cell nucleus was incorrectly calculated by multiplying the diploid genome size (in base pairs) by the mean weight of a nucleotide rather than of a nucleotide pair.

By using the data in Table 1, relative weights of nucleotide pairs can be calculated as follows: AT = 615.3830 and GC = 616.3711, bearing in mind that formation of

one phosphodiester linkage involves a loss of one H<sub>2</sub>O molecule. Further, phosphates of nucleotides in the DNA chain are acidic, so at physiologic pH the H<sup>+</sup> ion is dissociated (15). Provided the ratio of AT to GC pairs is 1:1, and ignoring the presence of modified nucleotides, the mean relative weight of one nucleotide pair is 615.8771. In any case, the error should be smaller than 1%.

The relative molecular weight may be converted to an absolute value by multiplying it by the atomic mass unit (1 u), which equals one-twelfth of a mass of <sup>12</sup>C, i.e.,  $1.660539 \times 10^{-27}$  kg (16). Consequently, the mean weight of one nucleotide pair would be  $1.023 \times 10^{-9}$  pg, and 1 pg of DNA would represent  $0.978 \times 10^9$  base pairs. The same conversion factor ( $0.98 \times 10^9$ ) was proposed by Cavalier-Smith (17), to our knowledge the only correct number provided in the relevant literature. In contrast, Straus (18) and Bennett and Smith (19) reported a lower value ( $0.965 \times 10^9$ ). The correct formulas for converting the number of nucleotide pairs to picograms of DNA and vice-versa are:

$$\text{genome size (bp)} = (0.978 \times 10^9) \times \text{DNA content (pg)}$$

$$\text{DNA content (pg)} = \text{genome size (bp)} / (0.978 \times 10^9)$$

The current estimates for human female and male diploid genome sizes are  $6.406 \times 10^9$  bp and  $6.294 \times 10^9$  bp, respectively (13). By using the conversion formulas given above, diploid human female and male nuclei in G<sub>1</sub> phase of the cell cycle should contain 6.550 and 6.436 pg of DNA, respectively. These values are in line with those of previous reports and differ drastically from those estimated by Thomas et al. (1).

Trout DNA content may be calculated by using the ratio of DNA amount of human versus trout. The ratio of 1.565 determined by Thomas et al. (1) seems too high when compared with other reports (12). The discrepancy might be due to the use of 4',6-diamidino-2-phenylindole to stain the samples, which binds preferentially to AT-rich DNA (20) and, hence, could result in the biased ratio of DNA amounts (11,21). Provided trout has 80% of human DNA content (12), trout nuclei should contain 5.149-5.240 pg

---

\*Correspondence to: Dr. J. Doležel, Laboratory of Molecular Cytogenetics and Cytometry, Institute of Experimental Botany, Sokolovská 6, CZ-77200 Olomouc, Czech Republic.

E-mail: dolez@ueb.cas.cz

Received 11 June 2001; Accepted 20 October 2002

Published online in Wiley InterScience (www.interscience.wiley.com).  
DOI: 10.1002/cyto.a.10013

Table 1  
Relative Molecular Weights of Nucleotides\*

Nucleotide	Chemical formula	Relative molecular weight
2'-deoxyadenosine 5'-monophosphate	C <sub>10</sub> H <sub>14</sub> N <sub>5</sub> O <sub>6</sub> P	331.2213
2'-deoxythymidine 5'-monophosphate	C <sub>10</sub> H <sub>15</sub> N <sub>2</sub> O <sub>8</sub> P	322.2079
2'-deoxyguanosine 5'-monophosphate	C <sub>10</sub> H <sub>14</sub> N <sub>5</sub> O <sub>7</sub> P	347.2207
2'-deoxycytidine 5'-monophosphate	C <sub>9</sub> H <sub>14</sub> N <sub>3</sub> O <sub>7</sub> P	307.1966

\*Calculated with the following standard atomic weights as provided by Vocke (14): A<sub>r</sub>(H) = 1.0079, A<sub>r</sub>(C) = 12.0107, A<sub>r</sub>(N) = 14.0067, A<sub>r</sub>(O) = 15.9994, A<sub>r</sub>(P) = 30.9738. Standard atomic weights are scaled to nuclide <sup>12</sup>C with A<sub>r</sub>(<sup>12</sup>C) = 12 and rounded to four decimals.

of DNA. These values agree with the published data and differ significantly from those reported by Thomas et al. (1).

#### LITERATURE CITED

- Thomas RA, Krishan A, Robinson DM, Sams C, Costa F. NASA/American Cancer Society High-Resolution Flow Cytometry Project—I. Cytometry 2001;43:2–11.
- Vinogradov AE. Genome size and GC-percent in vertebrates as determined by flow cytometry: the triangular relationship. Cytometry 1998;31:100–109.
- Vendrey R, Vendrey C. Arginine and deoxyribonucleic acid content of erythrocyte nuclei and sperms of some species of fishes. Nature 1953;172:30–31.
- Pedersen RA. DNA content, ribosomal gene multiplicity, and cell size in fish. J Exp Zool 1971;177:65–78.
- Tiersch TR, Chandler RW, Wachtel SSM, Ellias S. Reference standards for flow cytometry and application in comparative studies of nuclear DNA content. Cytometry 1989;10:706–710.
- Vendrey R, Vendrey C. La teneur du noyau cellulaire en acide désoxyribonucléique à travers les organes, les individus et les espèces animales. Experientia 1949;5:327–329.
- Mandel P, Métais P, Cuny S. Les quantités d'acide désoxypentose-nucléique par leucocyte chez diverses espèces de mammifères. C R Acad Sci Paris 1950;231:1172–1174.
- Métais P, Cuny S, Mandel P. La teneur en acide pentose-nucléique des leucocytes chez diverses espèces de mammifères et en particulier chez l'homme. C R Soc Biol 1951;145:1235–1238.
- Rasch EM. DNA "standards" and the range of accurate DNA estimates by Feulgen absorption microspectrophotometry. Adv Microsc 1985;196:137–166.
- Greilhuber J, Volleth M, Loidl J. Genome size of man and animals relative to the plant *Allium cepa*. Can J Genet Cytol 1983;25:554–560.
- Doležel J, Greilhuber J, Lucretti S, Meister A, Lysák MA, Nardi L, Obermayer R. Plant genome size estimation by flow cytometry: inter-laboratory comparison. Ann Bot 1998;82(suppl A):17–26.
- Vindeløv LL, Christensen IJ, Nissen NI. Standardization of high resolution flow cytometric DNA analysis by the simultaneous use of chicken and trout red blood cells as internal reference standards. Cytometry 1983;3:328–331.
- International Human Genome Sequencing Consortium. Initial sequencing and analysis of the human genome. Nature 2001;409:860–921.
- Vocke RD. Atomic weights of the elements 1997 (technical report). Pure Appl Chem 1999;71:1593–1607.
- Voet D, Voet JG, Pratt CW. Fundamentals of biochemistry. New York: John Wiley & Sons; 1999.
- Mohr PJ, Taylor BN. CODATA recommended values of the fundamental physical constants: 1998. J Phys Chem Ref Data 1999;28:1713–1852.
- Cavalier-Smith T. The evolution of genome size. New York: John Wiley & Sons; 1985.
- Straus NA. Comparative DNA renaturation kinetics in amphibians. Proc Natl Acad Sci USA 1971;68:799–802.
- Bennett MD, Smith JB. Nuclear DNA amounts in angiosperms. Phil Trans R Soc Lond B 1976;274:227–274.
- Portugal J, Waring MJ. Assignment of DNA binding sites for 4'-diamidino-2-phenylindole and bisbenzimidazole (Hoechst 33258). A comparative footprinting study. Biochim Biophys Acta 1988;949:158–168.
- Doležel J, Sgorbati S, Lucretti S. Comparison of three DNA fluorochromes for flow cytometric estimation of nuclear DNA content in plants. Physiol Plant 1992;85:625–631.

J. Doležel\*  
J. Bartoš

Laboratory of Molecular Cytogenetics and Cytometry  
Institute of Experimental Botany  
Olomouc, Czech Republic

H. Voglmayr  
J. Greilhuber  
Institute of Botany  
University of Vienna  
Vienna, Austria